(contents continued)

Severe learning deficits in apolipoprotein E-knockout mice in a water maze task
M.S. Oitzl, M. Mulder, P.J. Lucassen, L.M. Havekes, J. Grootendorst and E.R. De Kloet (The Netherlands) 189
Bicuculline administered into the amygdala after training blocks benzodiazepine-induced amnesia
H. Dickinson-Anson and J.L. McGaugh (USA) 197
The phospholipase A_2 inhibitor, quinacrine, reduces infarct size in rats after transient middle cerebral artery occlusion
A.Y. Estevez and J.W. Phillis (USA) 203
C57BL/6 strain is most susceptible to cerebral ischemia following bilateral common carotid occlusion among seven mouse strains: selective neuronal death in the murine transient forebrain ischemia
G. Yang, K. Kitagawa, K. Matsushita, T. Mabuchi, Y. Yagita, T. Yanagihara and M. Matsumoto (Japan) 209
C3H/HeJ mice are refractory to lipopolysaccharide in the brain
R.W. Johnson, G. Gheusi, S. Segreti, R. Dantzer and K.W. Kelley (USA, France) 219
Dopamine D_1 and D_2 receptor binding sites in brain samples from depressed suicides and controls
C. Bowden, A.E. Theodorou, S.C. Cheetham, S. Lowther, C.L.E. Katona, M.R. Crompton and R.W. Horton (UK) 227
Effects of ibogaine and noribogaine on the antinociceptive action of μ-, δ- and κ-opioid receptor agonists in mice
H.N. Bhargava, Y.-J. Cao and G.-M. Zhao (USA) 234
Inhibitory effects evoked from both the lateral and ventrolateral periaqueductal grey are selective for the nociceptive responses of rat dorsal horn neurones
A.J. Waters and B.M. Lumb (UK) 239
Effects of ibogaine on the development of tolerance to antinociceptive action of μ-, δ- and κ-opioid receptor agonists in mice
Y.-J. Cao and H.N. Bhargava (USA) 250
NMDA receptor dependence of the input specific NMDA receptor-independent LTP in the hippocampal CA1 region
M. Pananceau and B. Gustafsson (Sweden) 255
Predominant expression of Brn-2 in the postmitotic neurons of the developing mouse neocortex
K. Hagino-Yamagishi, Y. Saijoh, M. Ikeda, M. Ichikawa, R. Minamikawa-Tachino and H. Hamada (Japan) 261
High levels of human chymase expression in the pineal and pituitary glands
O. Baltatu, H. Nishimura, S. Hoffmann, G. Stoltenburg, I.D. Haulica, A. Lippoldt, D. Ganten and H. Urata (Germany, Romania, Japan) 269
Sex differences in bicuculline-induced convulsions: interaction with stress and ligands of benzodiazepine binding sites
D. Peričić and M. Bujas (Croatia) 279
Hypothyroidism coordinately and transiently affects myelin protein gene expression in most rat brain regions during postnatal development
N. Ibarrola and A. Rodríguez-Peña (Spain) 285
Protein synthesis-dependent and -independent mechanisms for the regulation of GnRH RNA transcript levels in GT1 cells
T.T. Yeo, A.C. Gore, M. Blum and J.L. Roberts (USA) 294
Increased calpain content and progressive degradation of neurofilament protein in spinal cord injury
N.L. Banik, D.C. Matzelle, G. Gantt-Wilford, A. Osborne and E.L. Hogan (USA) 301
Impact of pregnancy and lactation on $GABA_A$ receptor and central-type and peripheral-type benzodiazepine receptors
R. Weizman, E. Dagan, S.H. Snyder and M. Gavish (Israel) 307

Short communications

bFGF and FGFR-3 immunoreactivity in the rat brain following systemic kainic acid administration at convulsant doses: localization of bFGF and FGFR-3 in reactive astrocytes, and FGFR-3 in reactive microglia
J. Ballabriga, E. Pozas, A.M. Planas and I. Ferrer (Spain) 315
Lactation reduces Fos induction in the paraventricular and supraoptic nuclei of the hypothalamus after urethane administration in rats
B. Woodside and S. Amir (Canada) 319
Effect of lesion of the hypothalamic mamillary nuclei on skin blood vessel permeability in normal and capsaicin-pretreated rats
Z.S. Tolochko and G.B. Knyazeva (Russia) 324
DCG-IV, a potent metabotropic glutamate receptor agonist, as an NMDA receptor agonist in the rat cortical slice
Y. Uyama, M. Ishida and H. Shinozaki (Japan) 327

Author Index 331